

Exercise 1

By calculating the right and left derivatives of the following functions, determine which one is differentiable at a :

- 1 $f_1(x) = x^2 + |x + 1|$, $a = 1, -1$
- 2 $f_2(x) = \begin{cases} \frac{x}{1 + e^{1/x}}, & \text{si } x \in \mathbb{R}^* \\ 0, & \text{si } x = 0 \end{cases}, a = 0$

Exercise 2

Compute the derivatives of the following functions and precise their domains of definition.

- 1 $\sqrt[4]{x^3}$
- 2 $\frac{x}{x^3 + 1}$
- 3 $\frac{(1 + \sqrt{x})^3}{(x + 1)^2}$
- 4 $x \sqrt[n]{x}$, $n \in \mathbb{N}^*$
- 5 $x \ln |x + 1|$
- 6 $x^2 e^{1/x}$
- 7 $\sin(\cos(5x))$
- 8 a^x , $a \in \mathbb{R}^{+*}$
- 9 $(x + \ln x)^n$, $n \in \mathbb{N}^*$
- 10 $x^3 \ln(x)$
- 11 $x^2 e^x$

Exercise 3

Study the differentiability on \mathbb{R} of the following functions:

- 1 $f(x) = x|x|$
- 2 $g(x) = \frac{1}{2 + |x|}$
- 3 $h(x) = \begin{cases} x^2 \cos \frac{1}{x}, & \text{si } x \neq 0 \\ 0, & \text{si } x = 0 \end{cases}$

Exercice 4

Compute the n th derivative of the following functions

① $x\sqrt{x}$

② $\ln(x)$

③ e^{ax}

④ $\frac{1}{1-x}$

Exercice 5

Let a and b be two real numbers and f be a function defined on $[0, +\infty[$ by

$$f(x) = \begin{cases} 2\sqrt{x}, & \text{si } 0 \leq x \leq 1 \\ ax + b, & \text{si } x > 1 \end{cases}$$

Find a and b so that f is differentiable on $]0, +\infty[$

Exercice 6

Show that:

① $\forall x \in]0, \pi[: x \cos(x) - \sin(x) < 0$

② $\forall x \in]0, \frac{\pi}{2}[: \frac{2x}{\pi} < \sin(x) < x$

Exercice 7

In which of the following functions Rolle's theorem is applicable?

① $x^2 - 2$, on $[-2, 2]$

③ $\sqrt{1-x^2}$, on $[-1, 1]$

② $|x-2|$, on $[1, 3]$

④ $\tan(x)$, on $[\frac{\pi}{4}, \frac{\pi}{3}]$

Exercice 8

Let f be a function defined by

$$f(x) = e^{x^2} \cos(x)$$

Show that for all $a > 0$, the equation $f'(x) = 0$ has at least one solution on $[-a, a]$.

Exercise 9

- 1 apply the Mean value Theorem for the function $f : x \rightarrow x - x^3$ on the segment $[-2, 1]$ and compute the value $c \in]-2, 1[$ appearing in this formula.
- 2 apply the Mean value Theorem for the function $f : x \rightarrow x^2$ on the segment $[a, b]$ and compute the value $c \in]a, b[$ appearing in this formula.

Exercise 10

- 1 Using the Mean value Theorem, show that: $\frac{1}{1+x} < \ln(1+x) - \ln(x) < \frac{1}{x}$
- 2 Compute $\lim_{x \rightarrow +\infty} x[\ln(1+x) - \ln(x)]$
- 3 Deduce that: $\lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x}\right)^x = e$
- 4 Compute: $\lim_{x \rightarrow -\infty} \left(1 + \frac{1}{x}\right)^x$.