

Operators  
and  
Matrices

*With Compliments  
of the Author*

Zagreb, Croatia

Volume 13, Number 3, September 2019

*Ameur Seddik*

*Selfadjoint operators, normal operators, and  
characterizations*

# OPERATORS AND MATRICES

## AIMS AND SCOPE

'Operators and Matrices' (OaM), aims towards developing a high standard international journal which will publish top quality research papers in matrix and operator theory and their applications. The journal will publish mainly pure mathematics, but occasionally papers of a more applied nature could be accepted providing they have a nontrivial mathematical content. OaM will also publish expository papers and relevant book reviews.

OaM is published quarterly, in March, June, September and December.

## SUBMISSION

Manuscripts should be submitted through OaM page: [oam.ele-math.com](http://oam.ele-math.com), by e-mail of Editorial office, or directly to any one of the editors working in a related subject area.

Manuscripts are accepted for refereeing on the understanding that the same work has not been published (except in the form of an abstract), and is not under consideration for publication elsewhere.

Authors are advised to send notification of submission by e-mail to [oam@element.hr](mailto:oam@element.hr) (author(s), title, number of pages, name of editor to whom paper was sent to).

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send the file by electronic mail. The file must be the one from which the accompanying manuscript (finalized version) was printed out. A poscript or pdf file of this final version must also be enclosed.

## TITLE PAGE

The following data should be included on title page: the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author, and running head less than 60 characters.

## ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2010) and a list of 4-5 key words must be given.

## FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

## REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The authors should consult Mathematical Reviews for the standard abbreviations of journal names.

## PROOFS, PAGE CHARGES, OFFPRINTS

The galley proofs will be sent to corresponding author. Late return of the proofs will delay the article to a later issue.

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from OaM prior to publication.

## FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

OaM is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

[www.ele-math.com](http://www.ele-math.com)

e-mail: [oam@ele-math.com](mailto:oam@ele-math.com)

Operators and Matrices  
Editorial Office  
Menceticeva 2, 10000 Zagreb, Croatia  
Fax: +385 1 6008799

## SELFADJOINT OPERATORS, NORMAL OPERATORS, AND CHARACTERIZATIONS

AMEUR SEDDIK

(Communicated by F. Kittaneh)

*Abstract.* Let  $\mathfrak{B}(H)$  be the  $C^*$ -algebra of all bounded linear operators acting on a complex separable Hilbert space  $H$ . We shall show that:

1. The class of all selfadjoint operators in  $\mathfrak{B}(H)$  multiplied by scalars is characterized by

$$\forall X \in \mathfrak{B}(H), \|S^2X + XS^2\| \geq 2\|SXS\|, \quad (S \in \mathfrak{B}(H)).$$

2. The class of all normal operators in  $\mathfrak{B}(H)$  is characterized by each of the three following properties (where  $D_S = S^*S - SS^*$ , for  $S \in \mathfrak{B}(H)$ ),

- (i)  $\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq 2\|SXS\|, (S \in \mathfrak{B}(H)),$

- (ii)  $S^*D_S S = 0 = SD_S S^*, (S \in \mathfrak{B}(H)),$

- (iii)  $S^*D_S S \geq 0 \geq SD_S S^*, (S \in \mathfrak{B}(H)).$

### 1. Introduction and preliminaries

Let  $\mathfrak{B}(H)$  be the  $C^*$ -algebra of all bounded linear operators acting on a complex separable Hilbert space  $H$ .

We denote by:

- $D_S = S^*S - SS^*$ , the selfcommutator of  $S \in \mathfrak{B}(H)$ ,
- $(M)_1 = M \cap \{x \in X : \|x\| = 1\}$ , where  $X$  is any normed space and  $M$  is a subset of  $X$ ,
- $x \otimes y$  (where  $x, y \in H$ ), the operator (of rank less or equal to one) on  $H$  defined by  $(x \otimes y)z = \langle z, y \rangle x$ , for every  $z \in H$ .

It is known that:

- (i) for  $S \in \mathfrak{B}(H)$ ,  $S$  is with closed range if and only there exists a unique operator  $S^+$  (the Moore-Penrose inverse of  $S$ ) in  $\mathfrak{B}(H)$  such that  $SS^+S = S$ ,  $S^+SS^+ = S^+$ ,  $(SS^+)^* = SS^+$ , and  $(S^+S)^* = S^+S$ .
- (ii) if  $S \in \mathfrak{B}(H)$  and is invertible, then  $S^{-1} = S^+$ .

*Mathematics subject classification* (2010): 47A30, 47A05, 47B15.

*Keywords and phrases:* Closed range operator, Moore-Penrose inverse, group inverse, selfadjoint operator, unitary operator, normal operator, partial isometry operator, isometry operator, operator inequality.

**1.1. Selfadjoint operators and arithmetic-geometric-mean inequality**

In [4], Heinz proved that for every two positive operators  $P$  and  $Q$  in  $\mathfrak{B}(H)$ , and for every  $\alpha \in [0, 1]$ , the following operator inequality holds

$$\forall X \in \mathfrak{B}(H), \|PX + XQ\| \geq \|P^\alpha X Q^{1-\alpha} + P^{1-\alpha} X Q^\alpha\|. \tag{HI}$$

Making  $\alpha = \frac{1}{2}$  in the above inequality, we obtain the well known arithmetic-geometric mean inequality given by

$$\forall A, B, X \in \mathfrak{B}(H), \|A^*AX + XBB^*\| \geq 2\|AXB\|. \tag{AGMI1}$$

Note that the proof of (HI) given by Heinz is somewhat complicated. For this reason, McIntosh [5] with an elegant proof, proved that the operator inequality (AGMI1) holds, and deduced from it the Heinz inequality by iteration method.

Independently of the work of Heinz and McIntosh, Corach et al. proved in [1], that for every invertible selfadjoint operator  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|SXS^{-1} + S^{-1}XS\| \geq 2\|X\|, \tag{CPR1}$$

In [2], it was proved that the three above inequalities are equivalent, and we found an easy proof of (CPR1). This gives us another proof of Heinz inequality.

Recently [11], it was given three other inequalities equivalent to the Heinz inequality listed as follows:

- For every selfadjoint operator with closed range  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|SXS^+ + S^+XS\| \geq 2\|SS^+XS^+S\|. \tag{S1}$$

- For every selfadjoint operator with closed range  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|S^2X + XS^2\| \geq 2\|SXS\|. \tag{S2}$$

- For every selfadjoint operator  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|S^2X + XS^2\| \geq 2\|SXS\|. \tag{S3}$$

Then, we have a family of four operator inequalities generated by a selfadjoint operator (invertible, with closed range, and any) that are equivalent to the Heinz inequality (or to (AGMI1)), and each of them holds.

It is natural to know about the maximal form of each inequality of this family. This kind of problem was introduced in [6], where it was proved that an invertible operator  $S \in \mathfrak{B}(H)$  satisfies the inequality (CPR1) if and only if  $S$  is an invertible selfadjoint operator multiplied by a nonzero scalar. In [9, 10], it was shown that an operator with closed range  $S \in \mathfrak{B}(H)$  satisfies the inequality (S1) (or (S2)) if and only if  $S$  is a selfadjoint operator with closed range multiplied by a scalar. Note that this last

characterization concerning the operator with closed range case follows from the first characterization with the invertible operator case.

In this note, we shall show in the general situation, that an operator  $S \in \mathfrak{B}(H)$  satisfies the inequality (S3) if and only if  $S$  is a selfadjoint operator multiplied by a scalar.

### 1.2. Normal operators and arithmetic-geometric mean inequality

Recently [11], a second family of operator inequalities was given, a family of four inequalities that are equivalent to a second arithmetic-geometric mean inequality given by

$$\forall A, B, X \in \mathfrak{B}(H), \|A^*AX\| + \|XBB^*\| \geq 2\|AXB\|. \tag{AGMI2}$$

This family is given by:

- For every invertible normal operator  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|SXS^{-1}\| + \|S^{-1}XS\| \geq 2\|X\|. \tag{CPR12}$$

- For every normal operator with closed range  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|SXS^+\| + \|S^+XS\| \geq 2\|SS^+XS^+S\|. \tag{N1}$$

- For every normal operator with closed range  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq 2\|SXS\|. \tag{N2}$$

- For every normal operator  $S$  in  $\mathfrak{B}(H)$ , the following inequality holds

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq 2\|SXS\|. \tag{N3}$$

Note that the inequality (AGMI2) follows immediately from (AGMI1). In [11], a direct proof of (AGMI2) was given independently of (AGMI1).

As with the first above family, it is interesting to know about the maximal form of each inequality of this second family. In [7], it was proved that an invertible operator  $S \in \mathfrak{B}(H)$  satisfies the inequality (CPR12) if and only if  $S$  is an invertible normal operator. In [9, 10], it was shown that an operator with closed range  $S \in \mathfrak{B}(H)$  satisfies the inequality (N1) (or (N2)) if and only if  $S$  is a normal operator with closed range. Note that this last characterization concerning the operator with closed range case follows from the first characterization with the invertible operator case.

In this note, we shall show in the general situation, that an operator  $S \in \mathfrak{B}(H)$  satisfies the inequality (N3) if and only if  $S$  normal.

By using this last characterization in term of operator inequality of the class of normal operators, we shall show two general characterizations of this class given by

- $S^*D_S S = 0 = SD_S S^*$ , ( $S \in \mathfrak{B}(H)$ ).
- $S^*D_S S \geq 0 \geq SD_S S^*$ , ( $S \in \mathfrak{B}(H)$ ).

## 2. Characterizations of some distinguished classes of operators in terms of operator inequalities

In this section, we shall give characteristic properties in terms of operator inequalities for the two distinguished classes of operators, namely the class of all selfadjoint operators in  $\mathfrak{B}(H)$  multiplied by scalars, and the class of all normal operators in  $\mathfrak{B}(H)$ . Also, we shall present two general characterizations of this last class.

Along this section, let  $S \in \mathfrak{B}(H)$ , and since the class of all operators with closed ranges in  $\mathfrak{B}(H)$  is dense in  $\mathfrak{B}(H)$  (see [3, Ex. 140, p. 75]), we denote by  $(S_n)_{n \geq 1}$  a sequence of operators with closed ranges in  $\mathfrak{B}(H)$  that converges uniformly to  $S$ .

PROPOSITION 1. *The two following properties are equivalent:*

- (i) *S is a selfadjoint operator multiplied by a scalar,*
- (ii)  $\forall X \in \mathfrak{B}(H), \|S^2X + XS^2\| \geq 2\|SXS\|.$

*Proof.* We may assume without loss of generality that  $\|S\| = 1$ .

(i)  $\Rightarrow$  (ii). This implication follows immediately from (AGMI1).

(ii)  $\Rightarrow$  (i). Assume (ii) holds.

Applying triangular inequality in (ii), we deduce that  $\|S^2\| = \|S\|^2 = 1$ .

Define the real function  $F$  on the complete metric space  $(\mathfrak{B}(H))_1$  by  $F(X) = \|S^2X + XS^2\| - 2\|SXS\|$ , for  $X \in (\mathfrak{B}(H))_1$ ; and for  $n \geq 1$ , define the real function  $F_n$  on  $(\mathfrak{B}(H))_1$  by  $F_n(X) = \|S_n^2X + XS_n^2\| - 2\|S_nXS_n\|$ , for  $X \in (\mathfrak{B}(H))_1$ .

Put  $D = \{X \in (\mathfrak{B}(H))_1 : F(X) > 0\}$ . Then there are two cases,  $D = \emptyset, D \neq \emptyset$ .

Case 1.  $D = \emptyset$ . So, it follows that

$$\forall X \in \mathfrak{B}(H), \|S^2X + XS^2\| = 2\|SXS\|. \tag{*}$$

From this equality, we have

$$\forall x, y \in H, \|S^2x \otimes y + x \otimes S^{*2}y\| = 2\|Sx\| \|S^*y\|.$$

Using this last equality and since  $S^2 \neq 0$ , we deduce that  $\ker S^* = \{0\}$ . Hence,  $S$  is with dense range. Using again this last equality, we obtain the following inequality,

$$\forall x, y \in (H)_1, \|S^2x\| + 2\|Sx\| \|S^*y\| \geq \|S^{*2}y\|.$$

By taking the supremum over  $y \in (H)_1$ , we obtain that  $\|Sx\| \geq \frac{1}{3}\|x\|$ , for every  $x \in H$ . Thus,  $S$  is bounded below with dense range. Hence,  $S$  is invertible. So, from (\*), it follows that

$$\forall X \in \mathfrak{B}(H), \|SXS^{-1} + S^{-1}XS\| = 2\|X\|.$$

Then from [8],  $S$  is a unitary reflection operator in  $\mathfrak{B}(H)$  multiplied by a nonzero scalar. This gives us that (i) holds.

Case 2.  $D \neq \emptyset$ . From the fact that  $F$  is a positive continuous map on  $(\mathfrak{B}(H))_1$ , it follows that

$$\overline{D} = \overline{F^{-1}((0, \infty))} = F^{-1}([0, \infty)) = \{X \in (\mathfrak{B}(H))_1 : F(X) \geq 0\} = (\mathfrak{B}(H))_1.$$

Let  $X \in D$ , and  $\varepsilon > 0$ . Since  $S_n \rightarrow S$  uniformly, then there exists an integer  $N \geq 1$  (depends only in  $\varepsilon$ ) such that

$$\forall n \geq N, \forall Y \in (\mathfrak{B}(H))_1, |F(Y) - F_n(Y)| \leq \varepsilon.$$

If there exists  $n \geq N$  such that  $F_n(X) < 0$ , then using this last inequality, we have  $0 \leq F(X) < \varepsilon$ , for every  $\varepsilon > 0$ ; thus  $F(X) = 0$ , leading a contradiction with  $X \in D$ .

From this fact, it follows that

$$\forall X \in D, \forall n \geq N, F_n(X) \geq 0.$$

Since each  $F_n$  is a continuous map on  $(\mathfrak{B}(H))_1$  and  $D$  is dense in  $(\mathfrak{B}(H))_1$ , then

$$\forall X \in (\mathfrak{B}(H))_1, \forall n \geq N, F_n(X) \geq 0.$$

So, it follows that

$$\forall X \in \mathfrak{B}(H), \forall n \geq N, \|S_n^2 X + X S_n^2\| \geq 2 \|S_n X S_n\|.$$

Since for each  $n \geq 1$ ,  $S_n$  is with closed range, then from the corresponding characterization in the case of operators with closed ranges [9, 10], we obtain that  $S_n$  is a selfadjoint operator with closed range multiplied by a scalar, for every  $n \geq N$ . Since  $S_n \rightarrow S$  uniformly, and the class of all selfadjoint operators in  $\mathfrak{B}(H)$  is closed, then  $S$  is a selfadjoint operator multiplied by a scalar.  $\square$

REMARK 1. 1. The class of all selfadjoint operators in  $\mathfrak{B}(H)$  multiplied by scalars is characterized by

$$\forall X \in \mathfrak{B}(H), \|S^2 X + X S^2\| \geq 2 \|S X S\|, \quad (S \in \mathfrak{B}(H)).$$

2. From a part of the above proof, the class of all unitary reflection operators in  $\mathfrak{B}(H)$  multiplied by scalars is characterized by

$$\forall X \in \mathfrak{B}(H), \|S^2 X + X S^2\| = 2 \|S X S\|, \quad (S \in \mathfrak{B}(H)).$$

3. Proposition 1 gives us a positive answer of Problem 1 in [11].

PROPOSITION 2. *The two following properties are equivalent:*

(i)  $S$  is normal,

(ii)  $\forall X \in \mathfrak{B}(H), \|S^2 X\| + \|X S^2\| \geq 2 \|S X S\|.$

*Proof.* We may assume without loss of generality that  $\|S\| = 1$ .

(i)  $\Rightarrow$  (ii). Assume (i) holds and let  $X \in \mathfrak{B}(H)$ . From the fact that  $S$  is normal and applying (AGMI2), we deduce

$$\|S^2 X\| + \|X S^2\| = \|S^* S X\| + \|X S S^*\| \geq 2 \|S X S\|.$$

(ii)  $\Rightarrow$  (i). Assume (ii) holds.

Then  $\|S^2\| = \|S\|^2 = 1$ .

Define the real function  $G$  on the complete metric space  $(\mathfrak{B}(H))_1$  by  $G(X) = \|S^2X\| + \|XS^2\| - 2\|SXS\|$ , for  $X \in (\mathfrak{B}(H))_1$ ; and for  $n \geq 1$ , define the real function  $G_n$  on  $(\mathfrak{B}(H))_1$  by  $G_n(X) = \|S_n^2X\| + \|XS_n^2\| - 2\|S_nXS_n\|$ , for  $X \in (\mathfrak{B}(H))_1$ .

Put  $L = \{X \in (\mathfrak{B}(H))_1 : G(X) > 0\}$ . Then there are two cases,  $L = \emptyset$ ,  $L \neq \emptyset$ .

Case 1.  $L = \emptyset$ . So, it follows that

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| = 2\|SXS\|. \tag{*}$$

From this equality, we have

$$\forall x, y \in H, \|S^2x\| \|y\| + \|x\| \|S^{*2}y\| = 2\|Sx\| \|S^*y\|.$$

From this, and using the same argument as used in the same case of Proposition 1, we obtain that  $S$  is invertible.

So, from (\*), it follows that

$$\forall X \in \mathfrak{B}(H), \|SXS^{-1}\| + \|S^{-1}XS\| = 2\|X\|.$$

Then from [8],  $S$  is a unitary operator in  $\mathfrak{B}(H)$  multiplied by a nonzero scalar. This gives us that (i) holds.

Case 2.  $L \neq \emptyset$ . Using the same argument as used in the proof of Proposition 1 with the function  $F$ , we find  $\bar{L} = (\mathfrak{B}(H))_1$ , and there exists an integer  $N \geq 1$  such that

$$\forall X \in L, \forall n \geq N, G_n(X) \geq 0.$$

Since each  $G_n$  is a continuous map on  $(\mathfrak{B}(H))_1$  and  $L$  is dense in  $(\mathfrak{B}(H))_1$ , then

$$\forall X \in (\mathfrak{B}(H))_1, \forall n \geq N, G_n(X) \geq 0.$$

So, it follows that

$$\forall X \in \mathfrak{B}(H), \forall n \geq N, \|S_n^2X\| + \|XS_n^2\| \geq 2\|S_nXS_n\|.$$

Since for each  $n \geq 1$ ,  $S_n$  is with closed range, then from the corresponding characterization in the case of operators with closed ranges in [9, 10], we obtain that  $S_n$  is a normal operator, for every  $n \geq N$ . Since  $S_n \rightarrow S$  uniformly and the class of all normal operators in  $\mathfrak{B}(H)$  is closed, then  $S$  is a normal operator.  $\square$

REMARK 2. 1. The class of all normal operators in  $\mathfrak{B}(H)$  is characterized by

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq 2\|SXS\|, \quad (S \in \mathfrak{B}(H)).$$

2. From a part of the above proof, the class of all unitary operators in  $\mathfrak{B}(H)$  multiplied by scalars is characterized by

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| = 2\|SXS\|, \quad (S \in \mathfrak{B}(H)).$$

3. Proposition 2 gives us a positive answer of Problem 2 in [11].



### 3. Application: new general characterizations of the class of all normal operators

For  $S \in \mathfrak{B}(H)$ , from the definition,  $S$  is normal if and only if  $D_S = 0$ ; and it is well known that  $S$  is normal if and only if  $SD_S = D_S S$ . In this section, we shall present two new general forms of characterizations of the class of all normal operators.

PROPOSITION 3. *Let  $S \in \mathfrak{B}(H)$ . The following properties are equivalent:*

- (i)  $S$  is normal,
- (ii)  $S^*D_S S = 0 = SD_S S^*$ ,
- (iii)  $S^*D_S S \geq 0 \geq SD_S S^*$ .

*Proof.* The two implications (i)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (iii) are trivial.

(iii)  $\Rightarrow$  (i). Assume (iii) holds. So, we have

$$\begin{cases} \forall x \in H, \|S^2x\| \geq \|S^*Sx\|, \\ \forall x \in H, \|S^{*2}x\| \geq \|SS^*x\|. \end{cases}$$

Hence,

$$\begin{cases} \forall X \in \mathfrak{B}(H), \|S^2X\| \geq \|S^*SX\|, \\ \forall X \in \mathfrak{B}(H), \|XS^2\| \geq \|XSS^*\|. \end{cases}$$

So we obtain,

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq \|S^*SX\| + \|XSS^*\|.$$

Using (AGMI2), we deduce that

$$\forall X \in \mathfrak{B}(H), \|S^2X\| + \|XS^2\| \geq 2\|SXS\|.$$

Applying Theorem 2, (i) holds.  $\square$

REMARK 3. The class of all normal operators in  $\mathfrak{B}(H)$  is characterized by each of the two following properties:

- $S^*D_S S = 0 = SD_S S^*$ , ( $S \in \mathfrak{B}(H)$ ).
- $S^*D_S S \geq 0 \geq SD_S S^*$ , ( $S \in \mathfrak{B}(H)$ ).

*Acknowledgement.* The author would like to express his thanks to the referee for heart-warming suggestions.

## REFERENCES

- [1] G. CORACH, R. PORTA, AND L. RECHT, *An operator inequality*, Linear Algebra Appl., 142(1990), 153–158.
- [2] J. FUJII, M. FUJII, T. FURUTA, AND R. NAKAMOTO, *Norm inequalities equivalent to Heinz inequality*, Proc. Amer. Math. Soc. 118 (3) (1993), 827–830.
- [3] P. R. HALMOS, *A Hilbert space problem book*, Springer Second edition, 1982.
- [4] E. HEINZ, *Beiträge zur Störungstheorie der Spectralzerlegung*, Math. Ann. 123(1951), 415–438.
- [5] A. MCINTOSH, “*Heinz Inequalities and Perturbation of Spectral Families*”, Macquarie Mathematical Reports, Macquarie Univ., 1979.
- [6] A. SEDDIK, *Some results related to Corach-Porta-Recht inequality*, Proc. Amer. Math. Soc. 129(2001), 3009–3015.
- [7] A. SEDDIK, *On the injective norm and characterization of some subclasses of normal operators by inequalities or equalities*, J. Math. Anal. Appl. 351(2009), 277–284.
- [8] A. SEDDIK, *Characterization of the class of unitary operators by operator inequalities*, Linear and Multilinear algebra, 59(2011), 1069–1074.
- [9] A. SEDDIK, *Moore-Penrose inverse and operator inequalities*, Extracta mathematicae, 30(2015), 29–39.
- [10] A. SEDDIK, *Corrigendum to Moore-Penrose Inverse and Operator Inequalities*, Extracta Mathematicae, 30 (2015), 29–39, Extracta mathematicae, 32(2017), 209–211.
- [11] C. BOURAYA AND A. SEDDIK, *On the characterizations of some distinguished subclasses of Hilbert space operators*, Acta Sci. Math. (Szeged), 84 (2018), 611–627.

(Received March 3, 2019)

Ameur Seddik  
Department of Mathematics  
Faculty of Mathematics and Computer Science, University of Batna 2  
Batna, Algeria  
e-mail: a.seddik@univ-batna2.dz